

# Quantitative genetic (animal) model example in **R**

Gregor Gorjanc  
gregor.gorjanc@bfro.uni-lj.si

April 10, 2009

## Introduction

The following is just a quick introduction to quantitative genetic model, which is usually called animal model in animal breeding scenario. This model provides inferences on parameters such as genetic (additive/breeding, dominance, ...) values and possibly also co-variance components (additive genetic variance, heritability, ...). Very nice introduction to this topic is in Mrode (2005), which also gives a list of key references. We use example from this book and will therefore be very brief.

This note is mainly for educational purposes. There are quite some programs (e.g. Druet and Ducrocq (2006) mentions **ASReml**, **BGF90**, **DFREML**, **DMU**, **MATVEC**, **PEST/VCE** and **WOMBAT**) that can fit animal models in a general manner and we suggest to take a look at them instead of trying to reinvent the wheel in **R**.

In short animal model is an example of a mixed model:

$$\mathbf{y} = \mathbf{X}\mathbf{b} + \mathbf{Z}\mathbf{u} + \mathbf{e},$$

where  $\mathbf{y}$  represents a vector of observed (measured) phenotype values,  $\mathbf{b}$  and  $\mathbf{u}$  are vectors of unknown parameters for “fixed” and “random” effects, while  $\mathbf{X}$  and  $\mathbf{Z}$  are corresponding design matrices and finally  $\mathbf{e}$  is a vector of residuals. Assuming normal density for  $\mathbf{y}$  the following standard assumptions are taken:

$$\begin{pmatrix} \mathbf{y} \\ \mathbf{u} \\ \mathbf{e} \end{pmatrix} \sim N \begin{pmatrix} \mathbf{X}\mathbf{b} & \mathbf{V} & \mathbf{Z}\mathbf{G} & \mathbf{R} \\ \mathbf{0} & \mathbf{G}\mathbf{Z}' & \mathbf{G} & \mathbf{0} \\ \mathbf{0} & \mathbf{R} & \mathbf{0} & \mathbf{R} \end{pmatrix}, \mathbf{V} = \mathbf{Z}\mathbf{G}\mathbf{Z}' + \mathbf{R}$$

Up to now all this is as in usual mixed model. Genetic aspect comes from specification of covariance matrix between elements of  $\mathbf{u}$ , which usually represents sum of additive effects of genes of individuals in the pedigree. For a univariate model the covariance matrix of additive effect can be written as  $\mathbf{G} = \mathbf{A}\sigma_u^2$ , where  $\mathbf{A}$  is additive/numerator relationship matrix (Wright, 1922) and  $\sigma_u^2$  is additive genetic variance (Falconer and Mackay, 1996).

## Mixed model equations (MME)

Solution for  $\mathbf{b}$  i.e. (E)BLUE and  $\mathbf{u}$  i.e. (E)BLUP can be obtained from (Henderson, 1949; Goldberger, 1962; Henderson, 1963):

$$\hat{\mathbf{b}} = (\mathbf{X}\mathbf{V}^{-1}\mathbf{X})^{-1}\mathbf{X}\mathbf{V}^{-1}\mathbf{y},$$

and

$$\hat{\mathbf{u}} = \mathbf{G}\mathbf{Z}'\mathbf{V}^{-1}(\mathbf{y} - \mathbf{X}\hat{\mathbf{b}}),$$

but in a case with a lot of records the size of  $\mathbf{V}$  is huge and its direct inverse prohibitive if possible at all. Henderson (1950) presented the solution to this problem with so called mixed model equations:

$$\begin{pmatrix} \mathbf{X}'\mathbf{R}^{-1}\mathbf{X} & \mathbf{X}'\mathbf{R}^{-1}\mathbf{Z} \\ \mathbf{Z}'\mathbf{R}^{-1}\mathbf{X} & \mathbf{Z}'\mathbf{R}^{-1}\mathbf{Z} + \mathbf{G}^{-1} \end{pmatrix} \begin{pmatrix} \hat{\mathbf{b}} \\ \hat{\mathbf{u}} \end{pmatrix} = \begin{pmatrix} \mathbf{X}'\mathbf{R}^{-1}\mathbf{y} \\ \mathbf{Z}'\mathbf{R}^{-1}\mathbf{y} \end{pmatrix}.$$

## Data

We will use pedigree and data example from Mrode (2005). Example shows a beef breeding scenario with 8 individuals (animals), where 5 of them have phenotype records (pre-weaning gain in kg) and 3 three of them are without records and link others through the pedigree.

```
> library(GeneticsPed)
```

NOTE: THIS PACKAGE IS NOW OBSOLETE.

The R-Genetics project has developed an set of enhanced genetics packages to replace 'genetics'. Please visit the project homepage at <http://rgenetics.org> for informtion.

```
> data(Mrode3.1)
> (x <- Pedigree(x=Mrode3.1, subject="calf", ascendant=c("sire", "dam"),
+               ascendantSex=c("Male", "Female"), sex="sex"))
```

	calf	sex	sire	dam	pwg
1	S4	Male	S1	<NA>	4.5
2	S5	Female	S3	S2	2.9
3	S6	Female	S1	S2	3.9
4	S7	Male	S4	S5	3.5
5	S8	Male	S3	S6	5.0

## The model

For this baby BLUP example we will postulate the following model:

$$y_{ij} = s_i + a_j + e_{ij},$$

where  $y_{ij}$  is pre-weaning gain (kg) of calf  $j$  of sex  $j$ ;  $s_i$  are parameters of sex effect, while  $a_j$  are parameters of additive genetic effect for pre-weaning gain and finally  $e_{ij}$  is residual. Variances for  $a_j$  and  $e_{ij}$  are assumed as  $\mathbf{G} = \mathbf{A}\sigma_a^2$  with  $\sigma_a^2 = 20 \text{ kg}^2$  and  $\mathbf{R} = \mathbf{I}\sigma_e^2$  with  $\sigma_e^2 = 40 \text{ kg}^2$ .

## Setting up the MME

Observed/measured phenotype records:

```
> (y <- x$pwg)

[1] 4.5 2.9 3.9 3.5 5.0
```

Design matrix ( $\mathbf{X}$ ) for sex effect:

```
> X <- model.matrix(~ x$sex - 1)
> t(X)

      1 2 3 4 5
x$sexFemale 0 1 1 0 0
x$sexMale   1 0 0 1 1
attr("assign")
[1] 1 1
attr("contrasts")
attr(,"contrasts")$`x$sex`
[1] "contr.treatment"
```

Design matrix ( $\mathbf{Z}$ ) for additive genetic effect. Note that first three columns do not have indicators since these columns are for individuals without phenotype records and appear in the model only through the pedigree.

```
> (Z <- model.matrix(object=x, y=x$pwg, id=x$calf))

      S2 S1 S3 S4 S5 S6 S7 S8
1  0  0  0  0  1  0  0  0
2  0  0  0  0  0  1  0  0
3  0  0  0  0  0  0  1  0
4  0  0  0  0  0  0  0  1
5  0  0  0  0  0  0  0  1
```

Left hand side (LHS) of MME without  $\mathbf{G}^{-1}$ :

```
> LHS <- rbind(cbind(t(X) %*% X, t(X) %*% Z),
+             cbind(t(Z) %*% X, t(Z) %*% Z))
> ## or more efficiently
> (LHS <- rbind(cbind(crossprod(X), crossprod(X, Z)),
+             cbind(crossprod(Z, X), crossprod(Z))))
```

	x\$sexFemale	x\$sexMale	S2	S1	S3	S4	S5	S6	S7	S8
x\$sexFemale	2		0	0	0	0	1	1	0	0
x\$sexMale	0		3	0	0	0	1	0	0	1
S2	0		0	0	0	0	0	0	0	0
S1	0		0	0	0	0	0	0	0	0
S3	0		0	0	0	0	0	0	0	0
S4	0		1	0	0	0	1	0	0	0
S5	1		0	0	0	0	0	1	0	0
S6	1		0	0	0	0	0	0	1	0
S7	0		1	0	0	0	0	0	0	1
S8	0		1	0	0	0	0	0	0	1

and adding  $\mathbf{G}^{-1}$ , which is in this case  $\mathbf{A}^{-1}\alpha$  and  $\alpha = \frac{\sigma_e^2}{\sigma_a^2} = \frac{40}{20} = 2$ .

```
> ## We want Ainv for all individuals in the pedigree not only individuals
> ## with records
> x <- extend(x)
> Ainv <- inverseAdditive(x=x)
> sigma2a <- 20
> sigma2e <- 40
> alpha <- sigma2e / sigma2a
> q <- nIndividual(x)
> p <- nrow(LHS) - q
> (LHS[(p + 1):(p + q), (p + 1):(p + q)] <-
+ LHS[(p + 1):(p + q), (p + 1):(p + q)] + Ainv * alpha)
```

	S2		S1	S3		S4	S5	S6	S7	S8
S2	4	1.000000	1	0.000000	-2	-2	0	0		
S1	1	3.666667	0	-1.333333	0	-2	0	0		
S3	1	0.000000	4	0.000000	-2	1	0	-2		
S4	0	-1.333333	0	4.666667	1	0	-2	0		
S5	-2	0.000000	-2	1.000000	6	0	-2	0		
S6	-2	-2.000000	1	0.000000	0	6	0	-2		
S7	0	0.000000	0	-2.000000	-2	0	5	0		
S8	0	0.000000	-2	0.000000	0	-2	0	5		

Right hand side (RHS) of MME:

```

> RHS <- rbind(t(X) %*% y,
+             t(Z) %*% y)
> ## or more efficiently
> RHS <- rbind(crossprod(X, y),
+             crossprod(Z, y))
> t(RHS)

      x$sexFemale x$sexMale S2 S1 S3 S4 S5 S6 S7 S8
[1,]          6.8          13  0  0  0 4.5 2.9 3.9 3.5  5

```

## Solution

```

> sol <- solve(LHS) %*% RHS
> ## or more efficiently
> sol <- solve(LHS, RHS)
> t(sol)

      x$sexFemale x$sexMale      S2      S1      S3      S4      S5
[1,]      3.40443  4.358502 -0.0187701 0.09844458 -0.0410842 -0.008663123 -0.1857321
      S6      S7      S8
[1,] 0.1768721 -0.2494586 0.1826147

```

That's all folks! Well, all for the introduction. There are numerous issues covered in the literature. A good starting point is Mrode (2005) as already mentioned in the beginning.

## R Session information

```
> toLatex(sessionInfo())
```

- R version 2.8.1 (2008-12-22), i386-apple-darwin8.11.1
- Locale: C
- Base packages: base, datasets, grDevices, graphics, methods, stats, tools, utils
- Other packages: GeneticsPed 1.4.0, MASS 7.2-46, combinat 0.0-6, gdata 2.4.2, genetics 1.3.4, gtools 2.5.0-1, mvtnorm 0.9-5

## References

Druet, T. and Ducrocq, V. (2006). Innovations in software packages in quantitative genetics. In *8th World Congress on Genetics Applied to Livestock Production, Belo Horizonte, 2006-08-13/18*. Brazilian Society of Animal Breeding. Communication 27-10.

- Falconer, D. S. and Mackay, T. F. C. (1996). *Introduction to Quantitative Genetics*. Longman, Essex, U.K., 4th ed. edition.
- Goldberger, A. S. (1962). Best linear unbiased prediction in the generalized linear regression model. *J. Am. Stat. Assoc.*, 57(6):369–375.
- Henderson, C. R. (1949). Estimation of changes in herd environment. 32:709. (Abstract).
- Henderson, C. R. (1950). Estimation of genetic parameters. *Ann. Math. Stat.*, 9:309.
- Henderson, C. R. (1963). Selection index and expected genetic advance. In Hanson, W. D. and Robinson, H. F., editors, *Statistical genetics and plant breeding*, number 982, pages 141–163. National academy of sciences and national research council, Washington DC.
- Mrode, R. A. (2005). *Linear models for the prediction of animal breeding values*. CAB International, Wallingford, Oxon OX10 8DE, UK, 2 edition.
- Wright, S. (1922). Coefficients of inbreeding and relationship. *Am. Nat.*, 56:330–338.